

AD-A091 987

RHODE ISLAND UNIV KINGSTON DEPT OF MATHEMATICS  
OPTIMIZATION OF MULTICHANNEL PROCESSING.(U)  
SEP 80 O SHISHA

F/G 9/3

UNCLASSIFIED

AFOSR-77-3174  
AFOSR-TR-80-1146 NL

1 1 1 1  
A  
A091 987



END  
DATE  
FILMED  
1 81  
DTIC

AFOSR-TR- 80 - 1146

13  
me

LEVEL *44*

FINAL REPORT

ON

GRANT AFOSR-77-3174

OPTIMIZATION OF MULTICHANNEL PROCESSING

SEPTEMBER 1980

DTIC  
SELECTED  
NOV 13 1980  
C

DOC FILE COPY

80 11 06 101

Approved for public release;  
distribution unlimited.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER <b>AFOSR-TR-88-1146</b>	2. GOVT ACCESSION NO. <b>AD-A091</b>	3. RECIPIENT'S CATALOG NUMBER <b>987</b>	
4. TITLE (and Subtitle) <b>OPTIMIZATION OF MULTICHANNEL PROCESSING</b>		5. TYPE OF REPORT & PERIOD COVERED <b>FINAL / rept.</b>	
7. AUTHOR(s) <b>Professor Oved Shisha</b>		6. PERFORMING ORG. REPORT NUMBER <b>1</b>	
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>University of Rhode Island, Department of Mathematics, Kingston, R. I. 02881</b>		8. CONTRACT OR GRANT NUMBER(s) <b>AFOSR-77-3174</b>	
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F/2304/A3</b>	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>12-12</b>		12. REPORT DATE <b>SEP 1988</b>	
		13. NUMBER OF PAGES <b>10</b>	
		15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release; distribution unlimited.</b>			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  The subset selection algorithm is extended to search for a best subset from a large set of complex-valued basis functions. This algorithm is used to design digital finite-duration impulse response (FIR) filters having fewer coefficients than conventional FIR filters. An optimum conventional FIR filter is derived which has best uniform spacing of the fixed number of samples which are to be used, and examples are presented which show that, for the same number of coefficients, the complex-subset-selection filter can give			

SECRET  
better results than the optimum conventional filter.

The complex subset selection method is also applied to estimation of the frequencies of sinusoids in the presence of noise. A windowing technique is introduced to increase the efficiency and accuracy of the algorithm for frequency estimates. The results are compared with Cramer-Rao bounds.

↑

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

I. Introduction. During the contract period substantial progress was made on three basic problem areas:

- (1) The partial basis problem.
- (2) Least squares approximation with restricted range.
- (3) Mixed norm approximation.

In sections II, III, IV, we will discuss some of our work in each of these areas. In section V some other results are discussed.

II. The Partial Basis Problem. This problem was formulated by us in 1975 as a generalization of a problem of placing antenna elements optimally in a line array; independently, G. G. Lorentz studied a special case [11]. The problem has received considerable attention from researchers in recent years and elegant results have been obtained by us and others. This should prove to be a fruitful area for further investigation.

The partial basis problem can be stated in its general form as follows. Let  $X$  be a normed linear space, let  $f, h_0, \dots, h_{N-1}$  belong to  $X$  and let  $n$  be an integer,  $1 \leq n < N$ . For every sequence

$\mu = \{\mu_k\}_{k=1}^n$  of integers, with  $0 \leq \mu_1 < \dots < \mu_n \leq N-1$ , consider

$$e(\mu) = \min_{c_1, \dots, c_n} \left\| f - \sum_{k=1}^n c_k h_{\mu_k} \right\|.$$

The problem is to minimize  $e(\mu)$ ; it is of particular interest when  $X$  is one of the standard function spaces. The main results of [7] gives sufficient conditions for the "tail"  $h_{N-n}, \dots, h_{N-1}$  to be the unique best partial basis of size  $n$ . Typical theorems are:

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF DISSEMINATION TO LDC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer

Theorem 1. Let  $0 < a < b < \infty$  and let  $N, n$  be integers with  $1 \leq n < N$ .

Let  $f$  be a real function, continuous in  $[a, b]$  and assume that, for  $k = 0, 1, \dots, n$ ,  $(x^{k-N} f)^{(k)}$  exists and is  $\geq 0$  in  $(a, b)$ , with strict inequality there for  $k = n-1$  and  $k = n$ . Let  $0 \leq \mu_1 < \dots < \mu_n \leq N-1$  be integers such that  $\{\mu_1, \dots, \mu_n\} \neq \{N-n, \dots, N-1\}$ . Let  $1 \leq p \leq \infty$ .

Then

$$\min_{c_k \text{ real}} \left\| f(x) - \sum_{k=N-n}^{N-1} c_k x^k \right\|_{L^p(a,b)} <$$

$$\min_{c_k \text{ real}} \left\| f(x) - \sum_{k=1}^n c_k x^{\mu_k} \right\|_{L^p(a,b)}.$$

Theorem 2. Let  $0 \leq \alpha_{N-1} < \alpha_{N-2} < \dots < \alpha_0 < \frac{1}{2}$  and let  $1 \leq n < N$ ,  $n$  an integer. Let  $f$  be a real function with  $f^{(2k+1)}(0) = 0$ ,

$k = 0, \dots, N-1$ , and  $f^{(2k)}(x) > 0$  on  $(0, \pi]$  for  $k = 0, 1, \dots, N$ .

Assume  $f^{(2N-1)}(x)$  is continuous from the right at 0. Let

$0 \leq \mu_1 < \dots < \mu_n \leq N-1$  be integers such that  $\{\mu_1, \dots, \mu_n\} \neq \{N-n, \dots, N-1\}$ .

Let  $1 \leq p \leq \infty$ . Then

$$\min_{c_k \text{ real}} \left\| f(x) - \sum_{k=N-n}^{N-1} c_k \cos \alpha_k x \right\|_{L^p(0,\pi)} <$$

$$\min_{c_k \text{ real}} \left\| f(x) - \sum_{k=1}^n c_k \cos \alpha_{\mu_k} x \right\|_{L^p(0,\pi)}.$$

The theory of Tchebycheff systems is important for partial basis results and [7] contains contributions to this theory. A typical theorem:

Accession For	
NTIS GRA&I	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution	
Availability	
Dist	
Special	

Theorem 3. Let  $0 < \alpha_0 < \dots < \alpha_n$ . Then

$\cos \alpha_n x, \sin \alpha_0 x, -\cos \alpha_1 x, -\sin \alpha_1 x, \dots, (-1)^n \cos \alpha_n x, (-1)^n \sin \alpha_n x$

is an extended complete Tchebycheff system on  $[0, \pi]$  if and only if

$\alpha_n < 1/2$ .

Elegant results have been obtained by Lewis, Pinkus, and Shisha [9] giving sufficient conditions for the "head"  $h_0, \dots, h_{n-1}$  to be the unique best partial basis of size  $n$ . Perhaps the most striking result is this:

Theorem 4. Let  $0 < a < b < \infty$ ,  $1 \leq n < N$ ,  $f$  continuous and positive

on  $[a, b]$ ,  $(-1)^k f^{(k)}(x) > 0$  for  $a < x < b$ ,  $k = 1, \dots, n$  and

$1 \leq p \leq \infty$ . Then for  $s = 1, \dots, n+1$  any subsequence of length  $s$  of

$f, 1, x, \dots, x^{N-1}$  is a Tchebycheff system on  $[a, b]$  and

$$\min_{c_k \text{ real}} \left\| f(x) - \sum_{k=0}^{n-1} c_k x^k \right\|_{L^p(a, b)} < \min_{c_k \text{ real}} \left\| f(x) - \sum_{k=1}^n c_k x^{\mu_k} \right\|_{L^p(a, b)}$$

whenever  $0 \leq \mu_1 < \dots < \mu_n \leq N-1$  and  $\{\mu_1, \dots, \mu_n\} \neq \{0, 1, \dots, n-1\}$ .

The classical completely monotonic functions satisfy the hypotheses of Theorem 4 and hence are best approximated by  $1, x, \dots, x^{n-1}$ . Examples are  $f(x) = e^{-cx}$ ,  $c > 0$ ,  $[a, b] \subset (0, \infty)$  and  $f(x) = x^{-M}$ ,  $M = \text{positive integer}$ ,  $[a, b] \subset (0, \infty)$ .

The problem of finding the best partial basis computationally, for situations where theorems of the above type are inapplicable, was studied in [1]. An algorithm was presented, generalizing an algorithm of Hocking and Leslie [12], and a computer program was written and tested on some numerical examples. Also a theorem giving continuous dependence of a best approximation on the basis functions was given in [1]. This algorithm

was extended to complex-valued basis functions and used to design finite impulse response digital filters and to estimate frequencies of sinusoids in the presence of noise; cf. [4].

III. Least Squares Approximation with Restricted Range. This problem can be formulated as follow:

$$\begin{array}{l} \text{minimize} \\ h \in H \end{array} \int_a^b [f(x) - h(x)]^2 dx$$

subject to  $\ell(x) \leq p(x) \leq u(x)$  for all  $x$  in a set  $I$ . This includes the problem of positive approximation (where  $p(x) \geq 0$  is required) which has been studied by several investigators ([13], [14]).

The following characterization theorem has been obtained (under certain hypotheses on  $H$ ,  $f$ ,  $\ell$ ,  $u$ , and  $I$  which will not be specified here):

The approximation  $h^*$  in  $H$  is a solution of the least squares restricted range problem if and only if  $\ell \leq h^* \leq u$  on  $I$  and there exists points  $x_1, \dots, x_k$  in  $I$  and constants  $\sigma_1, \dots, \sigma_k$  such that:

- (i) For  $i = 1, \dots, k$ , either  $h^*(x_i) = \ell(x_i)$   
or  $h^*(x_i) = u(x_i)$
- (ii)  $\text{sign } \sigma_i = \begin{cases} +1 & \text{if } h^*(x_i) = u(x_i) \\ -1 & \text{if } h^*(x_i) = \ell(x_i) \end{cases}$
- (iii)  $\int_a^b (f-h^*)h_j = \sum_{i=1}^k \sigma_i h_j(t_i), \quad j = 1, \dots, n.$

(Here  $h_1, \dots, h_n$  is a basis for the set of approximations).



Computational algorithms of the Remes type have been formulated, convergence proofs obtained, and numerical examples solved in [6], [8].

IV. Mixed Norm Approximation. In designing beam patterns for antenna arrays and in designing the frequency response of a digital filter, one may wish to use different criteria of approximation for different intervals. For example, one might use a uniform norm

$$\max_{x \text{ in } I_1} |f(x) - p(x)|$$

in the passband interval  $I_1$  and a weighted least squares norm

$$\left( \int_{I_2} w(x) |f(x) - p(x)|^2 dx \right)^{1/2}$$

in the stopband interval  $I_2$ . Two possible approaches are:

$$(i) \quad \underset{p}{\text{minimize}} \quad \int_{I_2} w(x) |f(x) - p(x)|^2 dx$$

subject to  $-\epsilon \leq f(x) - p(x) \leq \epsilon$  for all  $x$  in  $I_1$  ;

and

$$(ii) \quad \text{minimize} \quad \left\{ \lambda \max_{x \text{ in } I_1} |f(x) - p(x)|^2 \right. \\ \left. + (1-\lambda) \int_{I_2} w(x) |f(x) - p(x)|^2 dx \right\}$$

where  $0 < \lambda < 1$ . The "mixed norm" problem (ii) can be converted to a quadratic programming problem, and solved; cf [2]. It is shown in [2] that, for  $f(x)$  of a certain type, (i) and (ii) are

equivalent in the following sense: If  $p_\lambda(x)$  solves the problem (ii) for some  $\lambda$ ,  $0 < \lambda < 1$ , then there exists an  $\epsilon > 0$  such that  $p_\lambda(x)$  solves problem (i). Conversely, if  $p_\epsilon(x)$  solves (i) for some  $\epsilon$ , there exists  $\lambda$ ,  $0 < \lambda < 1$  such that  $p_\epsilon(x)$  solves (ii). It is also shown that mixed norm approximation is related to vectorial approximation, developed by A. Bacopoulos and others [15]. In [6] the "mixed norm" in (ii) above is generalized to  $\lambda^r ||f-p||_1^r + (1-\lambda)^r ||f-p||_2^r$  where  $1 \leq r$  and  $|| \cdot ||_1, || \cdot ||_2$  are two specified norms. Again an equivalence theorem is obtained. The " $r = \infty$ " case, formulated as  $\max \{ \lambda ||f-p||_1, (1-\lambda) ||f-p||_2 \}$ , is studied carefully in [3] for  $||f-p||_1$  a discrete uniform norm and  $||f-p||_2$  a least square norm: a computational algorithm is formulated, proved to converge, and tested numerically. This algorithm is an analogue of an algorithm of C. L. Lawson, cf. [16].

V. Other Results. In this section we discuss other work partially supported by the Grant, namely [10] and [5]. Let  $-\infty < a < 0 < b < \infty$ , and  $f$  a real function, continuous on  $[a, b]$  but not a polynomial there. Given integers  $k, n$  denote

$$(1) \quad E_n(f) = \min \max_{a \leq x \leq b} |f(x) - p_n(x)|$$

where the minimum is taken over polynomials of degree at most  $n$ . Let  $E_n^k(f)$  denote the right hand side of (1) where the minimum is taken over all polynomials of degree at most  $n$  whose coefficient of  $x^k$  is 0. It follows from work in [17], [18] that if  $f^{(k)}$  exists and satisfies a Lipschitz condition of order  $\alpha$  ( $0 < \alpha < 1$ ) throughout  $[a, b]$  and if

$f^{(k)}(0) \neq 0$  then there exists a positive constant  $A_n$  such that

$$(2) \quad \frac{E_n^{(k)}(f)}{E_n(f)} \geq A_k n^\alpha, \quad n = 0, 1, \dots$$

In [10] a converse theorem has been obtained which essentially states that if inequality (2) holds, then  $f^{(k)}$  exists and satisfies a Lipschitz condition of order  $\alpha$  on each closed interval  $[a', b']$  with  $a < a' < b' < b$ . Hence a remarkable theorem is obtained identifying a

certain smoothness of  $f$  with the growth of  $\frac{E_n^{(k)}(f)}{E_n(f)}$ .

Note that this theorem could be considered a "partial basis" result of type described in Section II.

In conclusion we mention briefly [5] where Muntz-type closure theorems are obtained for sequences of the form

$$\left\{ w(t) t^{\alpha_k} \right\}_{k=1}^{\infty} \quad \text{on unbounded intervals.}$$

# References

1. Levasseur, K. M. and Lewis, J. T., "Antenna Design as a Partial Basis Problem", University of Rhode Island Mathematics Department Technical Report No. 66, 1977.
2. Cooper, W. C., Levasseur, K. M., and Shisha, O., "Design of Minimum Noise Digital Filters Using a Mixed Norm", University of Rhode Island Mathematics Department Technical Report No. 67, 1977.
3. Levasseur, K. M., "Computation of Best Mixed Norm Approximations", University of Rhode Island Mathematics Department Technical Report No. 72, 1978.
4. Khorrami, J. R., "Study of Digital Filter Design and Frequency Estimation by Complex Subset Selection", M. S. Thesis, University of Rhode Island, 1979.
5. Zalik, R. A., "Weighted Polynomial Approximation on Unbounded Intervals", Journal of Approximation Theory, 28 (1980), 113-119.
6. Levasseur, K. M., "Best Approximation with respect to Two Objectives", Ph.D. Thesis, University of Rhode Island, 1980.
7. Shisha, O., "Tchebycheff Systems and Best Partial Bases", to appear in Pacific Journal of Mathematics.
8. Levasseur, K. M. and Lewis, J. T., "Least Squares Approximation with Restricted Range", manuscript in preparation.
9. Lewis, J. T., Pinkus, A., and Shisha, O., manuscript on the partial basis problem, in preparation.
10. Hasson, M. and Shisha, O., Approximation by lacunary polynomials: a converse theorem, manuscript in preparation.
11. Lorentz, G. G., "Approximation by Incomplete Polynomials (problems and results)", in Padé and Rational Approximation - Theory and Applications, edited by E. B. Saff and R. S. Varga, Academic Press, New York, 1977, 289-302.
12. Hocking, R. P., and Leslie, R. N., "Selection of the Best Subset in Regression Analysis", Technometrics, 9 (1967), 531-540.
13. Deutsch, F., "Some Applications of Functional Analysis to Approximation Theory", Ph.D. Thesis, Brown Univ., 1965.
14. Mathews, J. W., "An Algorithm for Obtaining Best Approximations on Sets of Positive Polynomials", SIAM J. Num. Anal., 9 (1972), 734-742.

15. Bacopoulos, A., "Non-linear Chebyshev Approximation by Vector Norms", J. Approximation Theory, 2 (1969), 79-84.
16. Rice, J. R., The Approximation of Functions, Vol. 2, Advanced Topics, Addison-Wesley, Reading, Mass., 1969.
17. Hasson, M., "Derivatives of the Algebraic Polynomials of Best Approximation", J. Approximation Theory, 28 (1980).
18. Hasson, M., "Comparison between the Degrees of Approximation by Lacunary and Ordinary Algebraic Polynomials", J. Approximation Theory, 28 (1980).

END

DATE  
FILMED

1 - 81

DTIC